

One new and three newly recorded olethreutine moths (Lepidoptera, Tortricidae) from Japan

Yoshitsugu NASU

153-2, Nakado, Hashimoto, Wakayama, 648-0023 Japan; e-mail: fwik6205@mb.infoweb.ne.jp

Abstract *Rhopobota okui* Nasu, n. sp., is described from Japan. *Bactra* (*Chiloides*) *cerata* (Meyrick), *Eucosma lacteana* (Treitschke) and *Parepisimia catharota* (Meyrick), are newly recorded from Japan.

Key words *Bactra* (*Chiloides*) *cerata* (Meyrick), *Eucosma lacteana* (Treitschke), *Rhopobota okui* Nasu, n. sp., *Parepisimia catharota* (Meyrick), Tortricidae, Japan.

In the course of taxonomic study on Japanese olethreutine moths I have found one new species of *Rhopobota* and three hitherto unrecorded species of *Bactra*, *Eucosma* and *Parepisimia* from Japan. In the present paper I describe the new species and record the three species as new to the moth fauna of Japan, with illustrations of adults and genitalia.

The following acronyms are used for the locations of specimens: FK—Collection of Dr F. Komai, the Osaka University of Arts, Kanan-cho, Japan; TO—Collection of Dr T. Oku, Morioka, Japan; UOP—Collection of the Entomological Laboratory of Osaka Prefecture University, Sakai, Japan. Unless otherwise stated, all other specimens are in my collection.

***Bactra* (*Chiloides*) *cerata* (Meyrick) (Figs 1, 5, 9)**

Polychrosis cerata Meyrick, 1909, *J. Bombay nat. Hist. Soc.* **19**: 587.

Bactra cerata: Diakonoff, 1950: 285, pl. 6, fig. 27 (♂ genitalia), pl. 8, fig. 39 (♀ genitalia); Clarke, 1958: 307, pl. 152, figs 1, 1a (adult, ♂ genitalia); Clarke, 1976: 71, fig. 30 (♂ genitalia), pl. 6, fig. f (adult).

Bactra (*Chiloides*) *cerata*: Diakonoff, 1964: 47, figs 29, 30 (♂ ♀ genitalia); Diakonoff, 1973: 359; Diakonoff, 1982: 74; Kawabe, 1989: 43, fig. 49 (adult); Kawabe *et al.*, 1992: 105.

Chiloides cerata: Kuznetsov, 1988: 168, fig. 2–3 (♂ genitalia).


Bactra (*Chiloides*) *cerata insularis* Diakonoff, 1964, *Zool. Verh.* **70**: 49, fig. 34 (♀ genitalia).

Diagnosis. This species is a small moth with grayish brown ground color on the forewing (wing expanse 10–12 mm) and characterized by the three or four yellow-brown patches on base to 2/3 of wing (rarely obscure) (Fig. 1). The male and female genitalia are similar to those of some species of the subgenus, but differ in having a spine on the top of aedeagus and not having any cornuti on vesica (Fig. 5).

Material examined. JAPAN. Honshu: Wakayama Pref.: Fujinami, 1 ♂, 6. VIII. 1976 (F. Komai leg.), FK. Kyushu: Kagoshima Pref.: Yakushima I., Onoaida, 7 ♂ 7 ♀, 4–9. IX. 1979 (Y. Nasu leg.); Nakanoshima I., 1 ♀, 27–29. IV. 1990 (Y. Nasu leg.). Ryukyu: Okinawa Pref.: Okinawa I., Yona, 1 ♂, 15–18. V. 1998 (T. Ohno leg.). TAIWAN. Hualien Pref.: Hunyea Spa, 1 ♂, 31. V–2. VI. 1982 (T. Tanabe leg.), UOP.

Distribution. India (Assam), Sri Lanka, Thailand, Vietnam, Lesser Sunda Is, Fiji, Palau Is, West South New Guinea, Taiwan, Japan (Honshu, Kyushu, Ryukyu). New to Japan.

Host-plant. Unknown.



Figs 1-4. Adults. 1. *Bactra* (*Chiloides*) *cerata* (Meyrick), ♂. 2. *Eucosma lacteana* (Treitschke), ♂. 3. *Rhopobota okui* Nasu, n. sp., ♂, holotype. 4. *Parepisimia catharota* (Meyrick), ♂.

***Eucosma lacteana* (Treitschke) (Figs 2, 6, 10)**

Grapholitha lacteana Treitschke, 1835, *Schmett. Eur.* **10** (3): 113, 256.

Grapholitha (*Semasia*) *lacteana*: Heinemann, 1863: 169.

Grapholitha (*Paedisca*) *lacteana*: Wocke, 1871: 252.

Epiblema lacteana: Rebel, 1901: 116; Kennel, 1910: 281, pl. 86, fig. 18 (adult); Kennel, 1921: 558, pl. 21, fig. 27 (adult); Schütze, 1931: 189.

Eucosma (*Phaneta*) *lacteana*: Swatschek, 1958: 141; Hannemann, 1961: 132, fig. 260 (♂ genitalia), pl. 13, fig. 20 (adult); Leraut, 1980: 96.

Eucosma (*Eucosma*) *lacteana*: Obraztsov, 1968: 7.

Eucosma lacteana: Kuznetsov, 1985: 798; Kuznetsov & Jalava, 1988: 136; Emmet, 1988: 189; Emmet, 1991: 158; Razowski, 1996: 149; Leraut, 1997: 145.

Carpocapsa maritima Westwood & Humphreys, 1845, *Br. Moths* **2**: 138. [Synonymized by Kuznetsov, 1985: 798.]

Phaneta maritima: Pierce & Metcalfe, 1922: 70, pl. 24 (♂ ♀ genitalia).

Eucosma maritima: Meyrick, 1928: 551; Bradley, 1959: 71, pl. 8, fig. 89 (adult); Kuznetsov, 1975: 431; Kuznetsov, 1976: 104; Palm, 1982: 75; Palm, 1985: 74; Razowski, 1987: 79, pl. 6, fig. 6 (adult), fig. 52 (♂ genitalia), fig. 197 (♀ genitalia).

Eucosma (*Phaneta*) *maritima*: Swatschek, 1958: 142; Hannemann, 1961: 133, fig. 263 (♂ genitalia), pl. 15, fig. 23 (adult); Obraztsov, 1968: 15; Bentinck & Diakonoff, 1968: 115, pl. 17, fig. 4 (adult), pl. 68, figs 159a-c (♂ ♀ genitalia); Bradley *et al.*, 1972: 29; Bradley *et al.*, 1979: 9, 184, pl. 37, figs 1, 2 (adult); Leraut, 1980: 96.

Eucosma (*Calosetia*) *maritima*: Kuznetsov, 1978: 531, fig. 456-4 (♂ genitalia), fig. 457-2 (wing).

Semasia maritima: Benander, 1950: 119, pl. 7, fig. 33 (wing), fig. 11g (valva).

Grapholitha candidulana Nolken, 1870, *Arb. NatForschVer. Riga* (N. F.) **3**: 413.

Diagnosis. This species has a slender white forewing with two or three small dark-brown patches (wing expanse 12-14 mm) (Fig. 2). In general appearance, this moth is similar to *E.*

metzneriana (Treitschke, 1830), but differs from it in having a slender, shorter forewing, obscure dark-brown patches on the forewing, triangular uncus and narrower neck in the male genitalia (Fig. 6), and oblong lamella postvaginalis in the female genitalia (Fig. 10).

Material examined. JAPAN. Hokkaido: Koton, 5 ♂ 4 ♀, em. 1–14. VIII. 1960 (T. Oku leg.), reared from larvae on flowers and seed-heads of *Artemisia princeps*, TO, 2 ♂ 1 ♀, 4–7. VIII. 1961 (T. Oku leg.), 1 ♂, 8. X. 1961 (T. Oku leg.). NORWAY. Vang, 1 ♂, 28. VII. 1958 (P. K. Nielsen leg.) determined by P. K. Nielsen as *E. maritima*, UOP.

Distribution. Europe, Russia, Mongolia, Japan (Hokkaido). New to Japan.

Host-plant. Compositae: *Artemisia absinthium* L., *A. campestris* L., *A. maritima* L., *A. vulgaris* L. (Kennel, 1910, 1921; Schütze, 1931; Swatschek, 1958); *A. princeps* Pamp.


Biological note. The larvae spin together the flowers and seeds of *Artemisia* spp. According to Bradley *et al.* (1979), in September and October the larva feeds on the florets and unripe seeds of *A. maritima*, living in the flower spike and spinning the florets and leaves together to form a silken tube in the British Islands. When fully fed the larva leaves the spinning and constructs a cocoon amongst debris or in the earth, in which it overwinters before pupating. Pupation takes place in the cocoon in June. Adults fly in July and August.

***Rhopobota okui* Nasu, n. sp. (Figs 3, 7, 11)**

Diagnosis. This species is a small moth with olive-brown ground color on the forewing (wing expanse 9–12 mm) and characterized by a whitish median fascia and leaden white ocelloid patch (Fig. 3). The species is superficially similar to *R. kaempferiana* (Oku, 1971), but differs from it in having a shorter forewing, broader median fascia, whitish ocelloid patch (in *kaempferiana* leaden) on the forewing, larger uncus and socius, and a tuft of hooked thick setae on dorsal angle and a process on ventral angle of cucullus in the male genitalia (Fig. 7), and triangular ostium bursae in the female genitalia (Fig. 11).

Adult. Male (Fig. 3). Wing expanse 9–11 mm. Head olive-brown, face whitish. Antenna olive-brown. Labial palpus short, olive-brown, inner surface whitish; anterior end of the second segment and the third segment whitish. Thorax and tegula olive-brown. Forewing elongate, apex round, without costal fold. Ground color olive-brown. Costa olive-brown, with six pairs of whitish strigulae (costal strigulae) from apex to basal 1/3. Outward-oblique leaden streaks originating from each costal strigula; the first three streaks confluent, touching upper part of ocelloid patch; the fourth one gradually broadened, connected to inner lateral band of ocelloid patch. Basal patch indicated by some weak gray streaks, occupying basal 1/4 of wing. Median fascia consisting of two whitish bands, originating from the fifth and sixth costal strigulae, gradually broadened, reaching the middle of dorsum. Ocelloid patch leaden-white, margined laterally with leaden-white transversal bands, containing some dark brown dashes, with a dark brown spot on costal 1/3 of inner edge of the inner lateral band. Cilia brown, olivaceous on apex. Hindwing light grayish brown; cilia concolorous with wing.

Male genitalia (Fig. 7). Tegumen with a pair of large boot-shaped processes sublaterally (so-called unci). Socius large, finger-shaped, setose. Valva oblong, basal opening large, with a triangular process on middle of sacculus. Cucullus oblong, setose on inner surface, with a tuft of 12–14 hooked thick setae (two of them long) on dorsal angle and a long process on ventral angle. Aedeagus cone-shaped, with 19–22 deciduous cornuti.




Figs 5-6. Male genitalia. 5. *Bactra* (*Chiloides*) *cerata* (Meyrick), YN-892. 6. *Eucosma lacteana* (Treitschke), YN-74 (scales=0.5mm).

Female. Wing expanse 10-12 mm. Similar to male.

Female genitalia (Fig. 11). Papilla analis large, flat. Apophysis posterioris as long as apophysis anterioris. Ostium bursae triangular, located in an incision on posterior edge of sternite 7. Sternite 7 broadly sclerotized in square. Ductus bursae with a narrow sclerotized ring on posterior 1/3. Corpus bursae globular, with a large bifid sclerite on posterior part. Signa consisting of two small horn-shaped sclerites, located posteriorly.

Material examined. Holotype. ♂, JAPAN. Honshu: Nara Pref.: Mt Takatori-yama, 12. VII. 1991 (Y. Nasu leg.), genitalia slide YN-873, UOP. Paratypes. Honshu: Nara Pref.: Mt Takatori-yama, 1 ♂ 3 ♀, 12. VII. 1991 (Y. Nasu leg.), 1 ♀, 20. VII. 1991 (Y. Nasu leg.). Hyogo Pref.: Mt Futatabi-san, 1 ♀, em. 29. IX. 1976 (F. Komai leg.), reared from larva on fruit of *Ilex pedunculosa*, FK, 1 ♂ 1 ♀, em. 2. X. 1976 (F. Komai leg.), reared from larvae on fruits of *I. pedunculosa*, FK; Sanda, 1 ♂ 1 ♀, 16. VII. 1976 (F. Komai leg.), FK. Wakayama Pref.: Hashimoto, 1 ♂, 22. VI. 1990 (Y. Nasu leg.).



Figs 7-8. Male genitalia. 7. *Rhopobota okui* Nasu, n. sp., holotype, YN-873. 8. *Parepisimia catharota* (Meyrick), YN-898 (scales=0.5 mm).

Distribution. Japan (Honshu).

Host-plant. Aquifoliaceae: *Ilex pedunculosa* Miq.


Biological note. The larva feeds on the fruit of *Ilex pedunculosa*. Judging from the collecting data of adults, this moth may be bivoltine (adults: late June to July, late September to early October), and the larva may eat also the young leaves of the host instead of fruits in June.

Etymology. This species is named in honor of Dr T. Oku.

Parepisimia catharota (Meyrick) (Figs 4, 8)

Statherotis catharota Meyrick, 1928, *Exot. Microlepid.* 3: 443; Clarke, 1958: 595, pl. 296, figs 3-3a (adult, ♂ genitalia).

Parepisimia catharota: Diakonoff, 1975: 306 (as *catherota*, missp.); Kawabe, 1989: 61, fig. 95 (adult); Kawabe *et al.*, 1992: 108.



Figs 9-10. Female genitalia. 9. *Bactra* (*Chiloides*) *cerata* (Meyrick), YN-893. 10. *Eucosma lacteana* (Treitschke), YN-75 (scales=0.5 mm).

Diagnosis. This species has a blackish brown forewing with a large yellow-brown median fascia and triangular costal patch on apical 1/3 of costa (wing expanse 12 mm) (Fig. 4). This moth is superficially similar to *P. relapsa* (Meyrick, 1928), but differs from it in having the median fascia of same width throughout, smaller triangular costal patch before apex (in *relapsa* enlarged in large patch, to tornus), and valva with broader cucullus and costa without process in the male genitalia (Fig. 8). The species of *Parepisimia* sometimes have androconial or long hair-like scales at the costal area on upper surface in the male hindwing (Diakonoff, 1973), and this species has a long hair-pencil at that place.

Material examined. JAPAN. Ryukyu: Okinawa Pref.: Ishigaki I., Mt Omoto-dake, 1 ♂, 9. V. 1978 (Y. Arita leg.), 1 ♂, 8. V. 1998 (T. Ueda leg.); Itona, 1 ♂, 10. V. 1998 (T. Ueda leg.).

Distribution. Andaman Is, Thailand, Taiwan, Japan (Ryukyu). New to Japan.

Host-plant. Unknown.




Fig. 11. Female genitalia. 11. *Rhopobota okui* Nasu, n. sp., paratype, YN-874 (scale=0.5 mm).

Acknowledgments

I express my cordial thanks to Dr T. Oku, Morioka, Japan and Dr F. Komai, the Osaka University of Arts, Kanan-cho, Japan, for their kind advice and allowing me to examine the specimens. I am also obliged to Dr Y. Arita, the Meijo University, Nagoya, Japan, and Dr T. Ueda and Mr T. Ohno, UOP, for gift of materials, to Dr F. Komai for his assistance in photographing Tortricidae genitalia, to Dr H. Tanaka and Dr M. Shibao, the Osaka Prefectural Agricultural and Forestry Research Center, Habikino, Japan, and to Mr U. Jinbo, the Tokyo Metropolitan University, Hachioji, Japan, for their assistance in many aspects.

References

- Benander, P., 1950. Fjärilar. Lepidoptera. II. Småfjärilar, Microlepidoptera. Vecklarfjärilar (Tortricina). *Svensk Insektfauna* 10: 1-173, pls 1-9.
 Bentinck, G. A. G. and A. Diakonoff, 1968. De Nederlandse Bladrollers (Tortricidae). *Monografieën*

- ned. ent. Vereen* **3**: 1-201, pls 1-99.
- Bradley, J. D., 1959. An illustrated list of the British Tortricidae part 2: Olethreutinae. *Entomologist's Gaz.* **10**: 60-80, pls 1-19.
- Bradley, J. D., Fletcher, D. S. and P. E. S. Whalley, 1972. In Kloet, G. S. and W. D. Hincks (Eds), A check list of British insects (2nd edn). Part 2: Lepidoptera. *Handbk Ident. Br. Insects* **11** (2): 25-38.
- Bradley, J. D., Tremewan, W. G. and A. Smith, 1979. *British tortricoid Moths. Tortricidae: Olethreutinae*. viii, 336 pp., 43 pls. The Ray Society, London.
- Clarke, J. F. G., 1958. *Catalogue of the Type Specimens of Microlepidoptera in the British Museum (Natural History) described by Edward Meyrick* **3**. 600 pp., 298 pls. British Museum (Natural History), London.
- , 1976. Microlepidoptera: Tortricoidea. *Insects Micronesia* **9** (1): 1-144, 1-13 pls.
- Diakonoff, A., 1950. The type specimens of certain Oriental Eucosmidae and Carposinidae (Microlepidoptera) described by Edward Meyrick, together with descriptions of new Eucosmidae and Carposinidae in the British Museum (Natural History). *Bull. Br. Mus. nat. Hist. (Ent.)* **1**: 273-300, pls 3-8.
- , 1964. Further records and descriptions of the species of *Bactra* Stephens (Lepidoptera, Tortricidae). *Zool. Verh. Leiden* (70): 1-81, pl. 1.
- , 1973. The South Asiatic Olethreutini (Lepidoptera, Tortricidae). *Zool. Monogr. Rijismus. nat. Hist.* **1**. xxi, 700 pp. 15 pls. E. J. Brill, Leiden.
- , 1975. New Tortricoidea (Lepidoptera) from Southeast Asia in the British Museum (Natural History). *Zool. Meded. Leiden* **48**: 297-320, pls 1-8.
- , 1982. On a collection of some families of Microlepidoptera from Sri Lanka (Ceylon). *Zool. Verh. Leiden* (193): 1-124, pls 1-18.
- Emmet, A. M., 1988. Olethreutinae. In Emmet, A.M. (Ed.), *A Field Guide to the smaller British Lepidoptera* (2nd edn): 169-204. The British Entomological & Natural History Society, London.
- , 1991. Check showing the life history and habits of the British Lepidoptera. In Emmet, A.M. & J. Heath, *The Moths and Butterflies of the Great Britain and Ireland* **7** (2): 61-301. Haley Books, Colchester.
- Hannemann, H. J., 1961. Kleinschmetterlinge oder Microlepidoptera 1. Die Wickler (s. str.) (Tortricidae). *Tierwelt Dtl.* **48**: 1-233, pls 1-22.
- Heinemann, H. von, 1863. *Schmetterlinge Deutschlands und der Schweiz* (2) **1** (1). 248, 39 pp. C.A. Schwetschke & Shon, Braunschweig.
- Kawabe, A., 1989. Records and descriptions of the subfamily Olethreutinae (Lepidoptera: Tortricidae) from Thailand. *Microlepid. Thailand* (2): 23-82.
- Kawabe, A. and F. Komai, with J. Razowski, 1992. Tortricidae. In Heppner, J.B. & H. Inoue (Eds), Checklist. *Lepid. Taiwan* **1** (2): 103-109.
- Kennel, J. von, 1910. Tortricidae. In Spuler, A. (Ed.), *Die Schmetterlinge Europas* **2**: 238-296, **3**: pls 83-86. E. Schweizerbart'sche Verlagsbuch-handlung, Stuttgart.
- , 1921. Die palaarktischen Tortriciden. *Zoologica, Stuttg.* **21** (54): 545-742, pls 21-24.
- Kuznetzov, V. I., 1975. On the fauna of leaf-rollers (Lepidoptera, Tortricidae) of Mongolia. *Insects Mongolia* **3**: 408-437 (in Russian).
- , 1976. Leaf-rollers of the tribe Eucosmini (Lepidoptera, Tortricidae) of the southern part of the Far East. *Trudy zool. Inst. Leningr.* **62**: 70-108 (in Russian).
- , 1978. Tortricidae. In Medvedev, G. S. (Ed.), *Keys to the Insects of the European Part of the USSR* **4** (1): 193-680. Nauka Publishers, Leningrad. (In Russian).
- , 1985. Type specimens of tortricids of the supertribe Eucosmidii (Lepidoptera, Tortricidae) from collection of F. Treitschke in the Hungarian Museum of Natural History (Budapest). *Ent. Obozr.* **64**: 789-803 (in Russian).
- , 1988. [Leaf-rollers of the supertribe Gateclareanidii and Olethreutidii (Lepidoptera, Tortricidae) of the fauna of the North Vietnam]. *Syst. Insects Mites* **70**: 165-181 (in Russian).
- Kuznetzov, V. I. and J. Jalava, 1988. Soviet-Finnish entomological expeditions to southern Siberia 1982-1984. 2. Tortricidae (Lepidoptera). *Nota lepid.* **11**: 126-138.

- Leraut, P., 1980. *Liste systématique et synonymique des Lépidoptères de France, Belgique et Corse*. Supplément à Alexanor, Revue des Lépidoptéristes français et au Bulletin de la Société entomologique de France. 334 pp. Paris.
- , 1997. *Liste systématique et synonymique des Lépidoptères de France, Belgique et Corse* (2^e éd.). Supplément à Alexanor. 526 pp. Bergium.
- Meyrick, E., 1928 (reprinted in 1968). *A revised Handbook of British Lepidoptera*. 914 pp. E. W. Classey Ltd., Hampton.
- Obratzov, N. S., 1968. Die Gattungen der palaearktischen Tortricidae 2. Die Unterfamilie Olethreutinae 8. *Tijdschr. Ent.* **111**: 1-48, pls 1-11.
- Palm, E., 1982. Atlas over viklernes udbredelse i Danmark (Tortricidae & Cochylidae). *Dansk faun. Bibliik* **2**: 1-110.
- , 1985. Tortricidae. In Schnack, K. (Ed.), Katalog over de danske Sommerfugle. *Ent. Meddr* **52** (2-3): 67-76.
- Pierce, F. N. and J. W. Metcalfe, 1922 (reprinted in 1960). *The Genitalia of the Group Tortricidae of the Lepidoptera of the British Islands*. xxii, 101 pp., 34 pls. E. W. Classey, Feltham.
- Razowski, J., 1987. Motyle (Lepidoptera) Polski 7 (1) Eucosmini. *Monografie Fauny Pol.* **15**: 1-253, pls 1-15.
- , 1996. Tortricidae. In Karsholt, O. and J. Razowski (Eds), *The Lepidoptera of Europe. A distributional Checklist*: 130-157. Apollo Books, Stenstrup.
- Rebel, H., 1901. Famil. Pyralidae-Micropterigidae. In Staudinger, O. and H. Rebel, *Catalog der Lepidopteren des palaearktischen Faunengebietes* **2**: 1-368. R. Friedlander & Sohn, Berlin.
- Schütze, K. T., 1931. *Die Biologie der Kleinschmetterlinge unter besonderer Berücksichtigung ihrer Nährpflanzen und Erscheinungszeiten*. 235 pp. Verlag Internationalen Entomologischen Vereins E. V., Frankfurt am Main.
- Swatschek, B., 1958. Die Larvalsystematik der Wickler (Tortricidae und Carposinidae). *Abh. Larval-syst. Insekten* **3**: 1-269.
- Wocke, M., 1871. II. Microlepidoptera. In Staudinger, O. and M. Wocke, *Catalog der Lepidopteren des europaischen Faunengebietes*. xxxviii, 426 pp. Hofbuchhandlung von Hermann Budach, Dresden.

摘 要

日本産ヒメハマキガ (鱗翅目, ハマキガ科) の 1 新種と 3 新記録種 (那須義次)

Bactra (*Chiloides*) *cerata* (Meyrick) キモンヒメハマキ (新称)

前翅開張 10-12 mm. 前翅の地色は灰褐色, 基部から 2/3 の間に 3-4 個の黄褐色の斑紋がある (まれに不明瞭) ことで, 同属の他の種と識別は容易である. 分布: インド (アッサム), スリランカ, タイ, ベトナム, 小スンダ列島, フィジー, パラオ諸島, 西南ニューギニア, 台湾, 日本 (本州, 九州, 琉球). 日本新記録. 寄主植物: 不明.

Eucosma lacteana (Treitschke) ホソバシロヒメハマキ (新称)

前翅開張 12-14 mm. *E. metzneriana* (Treitschke) トビモンシロヒメハマキに外部表徴では類似するが, より前翅が細く, 小さいこと, 斑紋が不明瞭なこと, 雄交尾器の uncus が 3 角形であること, valva のくびれ部 (neck) が狭いこと, 雌交尾器の lamella postvaginalis が長方形であることで識別できる. 分布: ヨーロッパ, ロシア, モンゴル, 日本 (北海道). 日本新記録. 寄主植物: キク科: ヨモギ属の種. 日本ではヨモギの花序から飼育されている.

Rhopobota okui Nasu (新種) ソヨゴチビヒメハマキ (新称)

前翅開張 9-12 mm. 外部表徴では *R. kaempferiana* (Oku) ヤマツツジマダラヒメハマキに類似するが, より前翅が小さいこと, 中帯がより広いこと, 肛上紋が白っぽいことで識別できる. 雌雄交尾器での識別は容易である. 分布: 日本 (本州). 寄主植物: モチノキ科: ソヨゴ (果実).

Parepisimia catharota (Meyrick) ミナミキオビヒメハマキ (新称)

前翅開張 12 mm. 本種は近縁種の *P. relapsa* (Meyrick) に類似するが、中帯が同幅であること、前縁翅頂近くの三角紋が小さいこと、雄交尾器では幅広い cucullus を持つこと、valva の costa に突起を持たないことで識別できる。分布: アンダマン諸島, タイ, 台湾, 日本 (琉球). 日本新記録. 寄主植物: 不明.

(Accepted September 8, 1999)